

hep-ph/0610116
October 2006

Proton Decay in Teravolt Unification

P.H. Frampton

University of North Carolina, Chapel Hill, NC 27599-3255, USA

Abstract

Model building based on abelian quiver gauge theories gives models which resemble trinification, originally proposed with yottavolt unification. However, unification occurs in the teravolt range so proton decay must be completely excluded in the scalar sector. It is straightforward to accomplish this by a discrete symmetry which is a generalized baryon number B . Unlike in trinification, this is possible because quarks and leptons acquire masses from relevant operators which appear in four-dimensional conformal symmetry breaking.

One approach to the hierarchy, or naturalness, problem is to postulate conformality, four-dimensional conformal invariance at high energy, for the non gravitational extension of the standard model. The conformality approach suggested [1] in 1998 has made considerable progress. Models which contain the standard model fields have been constructed [3] and a model which grand unifies at about 4 TeV [4] has been examined. The latter is the subject of this note.

The original speculation [1] that such models may be conformal has been refined to exclude models which contain scalar fields transforming as adjoint representations because only if all scalars are in bifundamentals are there chiral fermions and, also only if all scalars are in bifundamentals, the one-loop quadratic divergences cancel in the scalar propagator. We regard it as encouraging that these two desirable properties select the same subset of models.

Another phenomenological encouragement stems from the observation [2] that the standard model representations for the chiral fermions can all be accommodated in bifundamentals of $SU(3)^3$ and can appear naturally in the conformality approach. The model building yields theories similar to the trinification proposed in [6]. There are three significant differences between trinification and conformality:

- Unification in conformality occurs in the TeV (Teravolt) range while in trinification it was in the $10^{15} GeV$ (Yottavolt) range.
- In trinification the gauge group was quasi-simple $SU(3) \times SU(3) \times SU(3) \times Z_3$ with one unique gauge coupling while in conformality the Z_3 is absent.
- Most importantly, the fermions in conformality acquire mass from relevant operators which appear in four-dimensional conformal symmetry breaking, not only from Yukawa coupling to the Higgs scalar.

In the present note we address the issue of proton decay. Under $SU(3)_C \times SU(3) \times SU(3)$, the quarks and leptons families each appear in the representations

$$[(3, 3^*, 1) + (1, 3, 3^*) + (3^*, 1, 3)] \quad (1)$$

for which the baryon numbers are respectively $B = +1/3, 0, -1/3$ for quarks, leptons and antiquarks.

The gauge bosons cannot transform quarks into leptons or vice versa because of the factoring out of the color $SU(3)_C$ group. So unlike in $SU(5)$ ([7]) proton decay is absent in the gauge sector.

The scalars are likewise in 27's according to

$$[(3, 3^*, 1) + (1, 3, 3^*) + (3^*, 1, 3)] + c.c. \quad (2)$$

although here, unlike for the fermions the complex conjugate representations must be included.

Fermion masses arise in trinification [6] from Yukawa couplings of the form

$$(3, 3^*, 1)_q (3^*, 1, 3)_q (1, 3, 3^*)_\phi \quad (3)$$

for the quarks and

$$(1, 3, 3^*)_f (1, 3, 3^*)_f (1, 3, 3^*)_\phi \quad (4)$$

for the leptons. Because these are two independent couplings, trinification has the feature of giving no relationship between quark and lepton masses.

There are additional Yukawa couplings possible which would violate baryon number B and cause catastrophically rapid proton decay with a TeV unification scale. Therefore these must be forbidden, as is achieved by assigning a generalization of baryon number $B = +1/3, 0 - 1/3$ respectively to the three representation listed.

$$(3, 3^*, 1)_\phi \quad B = 1/3 \quad (5)$$

$$(1, 3, 3^*)_\phi \quad B = 0 \quad (6)$$

$$(3^*, 1, 3)_\phi \quad B = -1/3 \quad (7)$$

Such assignments are very natural.

In phenomenological analysis of trinification [8], however, such a procedure is avoided in order to be able to obtain acceptable quark and lepton mass matrices. With teravolt unification such departure from the simplest case is not possible as proton decay must be totally forbidden.

However, the quark and lepton masses in conformality will acquire contributions in relevant operators $m_i \psi \psi$ from breaking of conformal symmetry. The pattern of these masses cannot yet be calculated, although a constraint on the pattern of the corresponding scalar masses has been suggested in [5] based on the cancellation of quadratic divergences.

In grand unification based on conformality, therefore, it is expected that proton decay will be completely absent in the non-gravitational theory. Gravitational effects may eventually destabilize the proton but with lifetime $\sim 10^{50}y$ far beyond any foreseeable experiment.

Equally or more important is the realization that consistency of the conformality with proton decay dictates that the quark and lepton masses receive significant contributions from the four-dimensional conformal symmetry breaking, not only from the Yukawa coupling to Higgs as universally assumed in previous studies of grand unification.

In the standard model, while the $Z^0, W^\pm$ masses are accurately predicted by a Higgs mechanism, there is no similar statement about fermion masses. My conclusion is that quark and lepton masses arise principally as relevant operators ^{#1} arising from breaking

^{#1}One important question is how such mass terms are induced above the electroweak scale. One possibility is via four fermion operators. I thank Edoardo Di Napoli for discussion.

of four-dimensional conformal invariance; a corollary is that the couplings of the standard Higgs scalar to quarks and leptons depart from the values usually assumed.

This work was supported in part by the U.S. Department of Energy under Grant No. DE-FG02-06ER-41418.

References

- [1] P.H. Frampton, Phys. Rev. **D60**, 041901 (1999). [hep-th/9812117](#). See also [2].
- [2] P.H. Frampton and C. Vafa. [hep-th/9903226](#).
- [3] P.H. Frampton, Phys. Rev. **D60**, 085004 (1999). [hep-th/9905042](#); *ibid* **D60**, 121901 (1999). [hep-th/9907051](#).
- [4] P.H. Frampton, Mod. Phys. Lett. **A18**, 1377 (2003). [hep-ph/0208044](#); P.H. Frampton, R.M. Rohm and T. Takahashi, Phys. Lett. **B567**, 265 (2003). [hep-ph/0302074](#).
- [5] E. Di Napoli and P.H. Frampton, Phys. Lett. **B638**, 374 (2006). [hep-th/0603065](#).
- [6] S.L. Glashow, in Proceedings of the Fifth Workshop on Grand Unification, Editors: K.Kang, H. Fried and P.H. Frampton, World Scientific (1984). pages 88-94.
- [7] H. Georgi and S.L. Glashow, Phys. Rev. Lett. **32**, 438 (1974); H. Georgi, H.R. Quinn and S. Weinberg, Phys. Rev. Lett. **33**, 451 (1974).
- [8] K.S. Babu, X.G. He and S. Pakvasa, Phys. Rev. **D33**, 763 (1986).